Clallam County Forage Fish

Spawn Data

- Spawn Site
- No Spawn Found
- Herring Holding Areas
- Herring Spawning Areas

Habitat Restoration Potential

- Very High (56 - 95)
- High (31 - 55)
- Moderate (11 - 30)
- Low (-14 to -10)
- Very Low (-55 to -15)

0 2.5 5 10 Kilometers

1:200,000

0 2.5 5 10 Miles

Waadah Island
Neah Bay
Clallam Bay
Pillar Pt
Observatory Pt
Angeles Pt
Ediz Hook
Port Angeles
Elwha River
Green Pt
Dungeness Spit
Sequim
Sequim Bay
Cape George

The Forage Fish Habitat Suitability Index provides an estimate of the suitability of a given site for forage fish spawning. The estimate is based on the quality of functional habitat and the degree of shoreline modification at that site. An index score of 30 or higher is very likely to support spawning habitat (Anchor 2001).

The forage fish spawn sample site data for Clallam County are a compilation of the North Olympic Salmon Coalition (NOSC) data points and the WDFW data from 1991 to 2007.

Maps created for NWStraits MRC Conference, Port Angeles, WA Nov. 2008 by Padilla Bay National Estuarine Research Reserve (SShull Sept. 2008)